

Stavudine (d4T, Zerit)

For additional information see Drugs@FDA:

<http://www.accessdata.fda.gov/scripts/cder/drugsatfda/index.cfm>

Formulations

Oral Solution: 1 mg/mL

Capsules: 15 mg, 20 mg, 30 mg, and 40 mg

Generic: d4T capsules and solution have been approved by the Food and Drug Administration (FDA) for manufacture and distribution in the United States.

Dosing Recommendations

Neonate/infant dose (birth to 13 days):

0.5 mg/kg twice daily.

Pediatric dose (14 days and body weight <30 kg):

1 mg/kg twice daily.

Adolescent (body weight ≥30 kg)/adult dose:

30 to <60 kg: 30 mg twice daily.

≥60 kg: 40 mg twice daily*.

* The World Health Organization (WHO) recommends 30 mg twice daily regardless of body weight in adults (see [Pediatric Use](#)).

Selected Adverse Events

- Mitochondrial toxicity
- Peripheral neuropathy
- Lipoatrophy
- Pancreatitis
- Lactic acidosis/severe hepatomegaly with hepatic steatosis (higher incidence than with other nucleoside reverse transcriptase inhibitors [NRTIs])
- Hyperlipidemia
- Insulin resistance/diabetes mellitus
- Rapidly progressive ascending neuromuscular weakness (rare)

Special Instructions

- d4T can be given without regard to food.
- Shake d4T oral solution well before use. Keep refrigerated; the solution will remain stable for 30 days.

Metabolism

- Renal excretion 50%. Decrease dose in renal dysfunction.

Drug Interactions (See also the [Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents](#)):

- *Renal elimination:* Drugs that decrease renal function could decrease stavudine clearance.
- *Other NRTIs:* Stavudine should not be administered in combination with zidovudine because of virologic antagonism.
- *Overlapping toxicities:* The combination of stavudine and didanosine is not recommended for initial therapy because of overlapping toxicities. Toxicities are more often reported in adults and include se-

rious, even fatal, cases of lactic acidosis with hepatic steatosis with or without pancreatitis in pregnant women.

- **Ribavirin and interferon:** Hepatic decompensation (sometimes fatal) has occurred in HIV/hepatitis C virus (HCV) coinfecting patients receiving combination antiretroviral therapy (cART), interferon, and ribavirin.

Major Toxicities:

- *More common:* Headache, gastrointestinal (GI) disturbances, skin rashes, hyperlipidemia, and fat maldistribution.
- *Less common (more severe):* Peripheral neuropathy and pancreatitis. Lactic acidosis and severe hepatomegaly with hepatic steatosis, including fatal cases, have been reported. The combination of stavudine with didanosine may result in enhanced toxicity (increased risk of fatal and nonfatal cases of lactic acidosis, pancreatitis, and hepatotoxicity), particularly in adults, including pregnant women. This combination should not be used for initial therapy.
- *Rare:* Increased liver enzymes, rapidly progressive ascending neuromuscular weakness.

Resistance: The International Antiviral Society-USA (IAS-USA) maintains a list of updated resistance mutations (see http://www.iasusa.org/resistance_mutations/index.html), and the Stanford University HIV Drug Resistance Database offers a discussion of each mutation (see <http://hivdb.stanford.edu/pages/GRIP/d4T.html>).

Pediatric Use: Although stavudine is FDA approved for use in children, its use is limited because it carries a higher risk of side effects associated with mitochondrial toxicity and a higher incidence of lipodystrophy than other NRTIs.

Data from multiple pediatric studies of stavudine alone or in combination with other antiretrovirals (ARVs) demonstrate that stavudine appears safe and is associated with clinical and virologic response¹⁻⁷. In resource-limited countries, stavudine is frequently a component of initial cART therapy with lamivudine and nevirapine in children, often as a component of fixed-dose combinations not available in the United States. In this setting, reported outcomes from observational studies are good; data show substantial increases in the CD4 count and complete viral suppression in 50%–80% of treatment-naïve children⁸⁻¹¹. In such a setting, where pediatric patients are already predisposed to anemia because of malnutrition, parasitic infestations, or sickle cell anemia, stavudine carries a lower risk of hematologic toxicity than zidovudine, especially in those patients requiring cotrimoxazole prophylaxis¹².

Stavudine is associated with a higher rate of adverse events than zidovudine in adults and children receiving cART¹³⁻¹⁴. In a large pediatric natural history study (PACTG 219C), stavudine-containing regimens had a modest but significantly higher rate of clinical and laboratory toxicities than those containing zidovudine, with pancreatitis, peripheral neuropathy, and lipodystrophy/lipodystrophy (fat maldistribution) associated more often with stavudine use¹⁴. Peripheral neuropathy is an important toxicity associated with stavudine but appears to be less common in children than in adults^{2, 15}. In PACTG 219C, peripheral neuropathy was recognized in 0.9% of children¹⁴. Lipodystrophy, and specifically lipodystrophy (loss of subcutaneous fat), are toxicities associated with the use of NRTIs, particularly stavudine, in adults and children¹⁶⁻¹⁹. Lipodystrophy developed in 28% of 39 children receiving stavudine, lamivudine, and nelfinavir after a median of 49 months of therapy, with 9 children demonstrating lipodystrophy²⁰. Among 90 children receiving stavudine, lamivudine, and either nevirapine or efavirenz, 65% developed lipodystrophy at 33 months²¹.

Lactic acidosis with hepatic steatosis, including fatal cases, has been reported with the use of nucleoside analogues, including stavudine, alone or in combination²²⁻²⁴. The combination of stavudine and didanosine in pregnant women has been associated with fatal lactic acidosis and should be used during pregnancy only if no other alternatives are available²⁵. (For additional information on lactic acidosis see [Table 17g Antiretroviral Therapy-Associated Adverse Effects and Management Recommendations](#).)

Many of the above-mentioned adverse events are believed to be due to mitochondrial toxicity resulting from inhibition of mitochondrial DNA polymerase gamma, with depletion of mitochondrial DNA in fat, muscle, peripheral blood mononuclear cells (PBMCs), and other tissues^{22, 26-28}. In a recent analysis involving a large cohort of pediatric patients (Pediatric AIDS Clinical Trials Group protocols 219 and 219 C), possible mitochondrial dysfunction was associated with NRTI use, especially in those children receiving stavudine and/or lamivudine²⁹.

WHO recommends that stavudine be phased out of use because of serious, irreversible side effects and that a maximum stavudine dose of 30 mg be used instead of the FDA-recommended 40 mg in adults weighing 60 kg or more. Several studies have compared the efficacy and toxicity of the two doses: HIV suppression was found to be similar in adult patients treated in South Africa with either the 30-mg or 40-mg dose³⁰; the incidence of peripheral neuropathy in adults treated in South Africa was significantly lower in the 30-mg group than in the 40-mg group, but the overall incidence was considered to be unacceptably high³¹. To reduce the risk of or to manage toxicity, some Panel members support switching to another agent if available rather than lowering the maximum dose. This recommendation is based on the availability of alternative ARV agents in the United States and on concerns for underdosing some patients with stavudine. However, other Panel members prefer using the 30-mg maximum dose of stavudine when there are limited alternatives.

References

1. Aboulker JP, Babiker A, Chaix ML, et al. Highly active antiretroviral therapy started in infants under 3 months of age: 72-week follow-up for CD4 cell count, viral load and drug resistance outcome. *AIDS*. 2004;18(2):237-245.
2. Kline MW, Dunkle LM, Church JA, et al. A phase I/II evaluation of stavudine (d4T) in children with human immunodeficiency virus infection. *Pediatrics*. 1995;96(2 Pt 1 (Aug)):247-252.
3. Kline MW, Fletcher CV, Federici ME, et al. Combination therapy with stavudine and didanosine in children with advanced human immunodeficiency virus infection: pharmacokinetic properties, safety, and immunologic and virologic effects. *Pediatrics*. 1996;97(6 Pt 1):886-890.
4. Kline MW, van Dyke RB, Lindsey J, et al. Combination therapy with stavudine (d4T) plus didanosine (ddI) in children with human immunodeficiency virus infection. The Pediatric AIDS Clinical Trials Group 327 Team. *Pediatrics*. 1999;103(5):e62.
5. Krogstad P, Lee S, Johnson G, et al. Nucleoside-analogue reverse-transcriptase inhibitors plus nevirapine, nelfinavir, or zidovudine for pretreated children infected with human immunodeficiency virus type 1. *Clin Infect Dis*. 2002;34(7):991-1001.
6. Nachman SA, Stanley K, Yogev R, et al. Nucleoside analogs plus zidovudine in stable antiretroviral therapy-experienced HIV-infected children: a randomized controlled trial. Pediatric AIDS Clinical Trials Group 338 Study Team. *JAMA*. 2000;283(4):492-498.
7. Yogev R, Lee S, Wiznia A, et al. for the Pediatric AIDS Clinical Trials Group 338 Study Team. Stavudine, nevirapine and zidovudine in stable antiretroviral therapy-experienced children with human immunodeficiency virus infection. *Pediatr Infect Dis J*. 2002;21(2):119-125.

8. Bolton-Moore C, Mubiana-Mbewe M, Cantrell RA, et al. Clinical outcomes and CD4 cell response in children receiving antiretroviral therapy at primary health care facilities in Zambia. *JAMA*. 2007;298(16):1888-1899.
9. Janssens B, Raleigh B, Soeung S, et al. Effectiveness of highly active antiretroviral therapy in HIV-positive children: evaluation at 12 months in a routine program in Cambodia. *Pediatrics*. 2007;120(5):e1134-1140.
10. Kanya MR, Mayanja-Kizza H, Kambugu A, et al. Predictors of long-term viral failure among ugandan children and adults treated with antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2007;46(2):187-193.
11. Zhang F, Haberer JE, Zhao Y, et al. Chinese pediatric highly active antiretroviral therapy observational cohort: a 1-year analysis of clinical, immunologic, and virologic outcomes. *J Acquir Immune Defic Syndr*. 2007;46(5):594-598.
12. Okechukwu AA, Gambo D, Okechukwu IO. Prevalence of anaemia in HIV-infected children at the University of Abuja Teaching Hospital, Gwagwalada. *Niger J Med*. 2010;19(1):50-57.
13. Robbins GK, De Gruttola V, Shafer RW, et al. Comparison of sequential three-drug regimens as initial therapy for HIV-1 infection. *N Engl J Med*. 2003;349(24):2293-2303.
14. Van Dyke RB, Wang L, Williams PL. Toxicities associated with dual nucleoside reverse-transcriptase inhibitor regimens in HIV-infected children. *J Infect Dis*. 2008;198(11):1599-1608.
15. Kline MW, Fletcher CV, Harris AT, et al. A pilot study of combination therapy with indinavir, stavudine (d4T), and didanosine (ddI) in children infected with the human immunodeficiency virus. *J Pediatr*. 1998;132(3 Pt 1):543-546.
16. Joly V, Flandre P, Meiffredy V, et al. Increased risk of lipodystrophy under stavudine in HIV-1-infected patients: results of a substudy from a comparative trial. *AIDS*. 2002;16(18):2447-2454.
17. European Paediatric Lipodystrophy Group. Antiretroviral therapy, fat redistribution and hyperlipidaemia in HIV-infected children in Europe. *AIDS*. 2004;18(10):1443-1451.
18. Ene L, Goetghebuer T, Hainaut M, et al. Prevalence of lipodystrophy in HIV-infected children: a cross-sectional study. *Eur J Pediatr*. 2007;166(1):13-21.
19. Haubrich RH, Riddler SA, DiRienzo AG, et al. Metabolic outcomes in a randomized trial of nucleoside, nonnucleoside and protease inhibitor-sparing regimens for initial HIV treatment. *AIDS*. 2009;23(9):1109-1118.
20. Scherpbier HJ, Bekker V, van Leth F, et al. Long-term experience with combination antiretroviral therapy that contains nevirapine for up to 7 years in a pediatric cohort. *Pediatrics*. 2006;117(3):e528-536.
21. Aupibul L, Puthanakit T, Lee B, et al. Lipodystrophy and metabolic changes in HIV-infected children on non-nucleoside reverse transcriptase inhibitor-based antiretroviral therapy. *Antivir Ther*. 2007;12(8):1247-1254.
22. Haugaard SB, Andersen O, Pedersen SB, et al. Depleted skeletal muscle mitochondrial DNA, hyperlactatemia, and decreased oxidative capacity in HIV-infected patients on highly active antiretroviral therapy. *J Med Virol*. 2005;77(1):29-38.
23. Koh MT. Unrecognized near-fatal hyperlactatemia in an HIV-infected infant exposed to nucleoside reverse transcriptase inhibitors. *Int J Infect Dis*. 2007;11(1):85-86.
24. Hernandez Perez E, Dawood H. Stavudine-induced hyperlactatemia/lactic acidosis at a tertiary communicable diseases clinic in South Africa. *J Int Assoc Physicians AIDS Care (Chic)*. 2010;9(2):109-112.
25. Sarner L, Fakoya A. Acute onset lactic acidosis and pancreatitis in the third trimester of pregnancy in HIV-1 positive women taking antiretroviral medication. *Sex Transm Infect*. 2002;78(1):58-59.
26. Blanco F, Garcia-Benayas T, Jose de la Cruz J, et al. First-line therapy and mitochondrial damage: different nucleosides, different findings. *HIV Clin Trials*. 2003;4(1):11-19.
27. Cherry CL, Gahan ME, McArthur JC, et al. Exposure to dideoxynucleosides is reflected in lowered mitochondrial DNA in subcutaneous fat. *J Acquir Immune Defic Syndr*. 2002;30(3):271-277.

28. Sanchez-Conde M, de Mendoza C, Jimenez-Nacher I, et al. Reductions in stavudine dose might ameliorate mitochondrial-associated complications without compromising antiviral activity. *HIV Clin Trials*. 2005;6(4):197-202.
29. Crain MJ, Chernoff MC, Oleske JM, et al. Possible mitochondrial dysfunction and its association with antiretroviral therapy use in children perinatally infected with HIV. *J Infect Dis*. 2010;202(2):291-301.
30. Hoffmann CJ, Charalambous S, Fielding KL, et al. HIV suppression with stavudine 30 mg versus 40 mg in adults over 60 kg on antiretroviral therapy in South Africa. *AIDS*. 2009;23(13):1784-1786.
31. Pahuja M, Glesby MJ, Grobler A, et al. Effects of a reduced dose of stavudine (d4T) on the incidence and severity of peripheral neuropathy in PLHIV in South Africa. Paper presented at IAS-AIDS 2010; July 18–23, 2010; Vienna, Austria.